



Review

Pathophysiologic changes due to TASER® devices versus excited delirium: Potential relevance to deaths-in-custody?

James R. Jauchem, Ph.D, Senior Research Physiologist*

U.S. Air Force Research Laboratory, 711th Human Performance Wing, Human Effectiveness Directorate, Directed Energy Bioeffects Division, 8262 Hawks Road, Brooks City-Base, Texas 78235, USA

ARTICLE INFO

Article history:

Received 4 January 2010
Received in revised form
13 January 2011
Accepted 26 January 2011
Available online 18 February 2011

Keywords:

Electronic control device
Electric shocking device
TASER
Excited delirium
Electric stimulation
Conducted energy weapon

ABSTRACT

The syndrome of excited delirium has been implicated in some deaths-in-custody which also involved the use of electronic control devices (ECDs) (including those manufactured by TASER International) on subjects. This review is an update on recent studies of pathophysiologic changes related to these two separate but parallel topics: a) first, the use of ECDs during law-enforcement activities; and b) second, the occurrence of excited delirium during such activities. This is a narrative review of elements that may be of use in generating hypotheses relating to potential similarities or differences between the two topics. Differences between changes in most factors due to excited delirium versus those of ECD applications were not readily apparent in most cases. These factors include: direct and indirect effects on the cardiovascular system, respiration, rhabdomyolysis and muscle enzymes, hyperkalemia, acidosis, hyperglycemia, and increased hematocrit. One factor that may exhibit consistent differences, however, is increased body temperature, which is often evident during excited delirium (versus a lack of increase temperature during ECD exposures). Thus, on the basis of this review, a more detailed delineation of this factor could be a major focus for future forensic investigations of deaths-in-custody involving either excited delirium or ECD exposures.

© 2011 Elsevier Ltd and Faculty of Forensic and Legal Medicine. All rights reserved.

1. Introduction

“Electronic control devices” (ECDs) (such as those manufactured by TASER International [Scottsdale, Arizona, USA]) have been implicated in some deaths-in-custody (reviewed previously in this journal).¹ The syndrome referred to as “excited delirium,” however, is one possible explanation for some of these deaths. Since that first review in this journal (completed three years ago), additional research studies, using both animals and human subjects, have been completed in several laboratories. The purpose of the current review is to provide an update on the two parallel topics ([a] the use of ECDs, and [b] the occurrence of excited delirium), and to provide additional insights into possible improvements in pathological examinations of these deaths. This is a narrative review of elements that may be of use in generating hypotheses relating to potential similarities or differences between the two topics.

Because extreme applications of ECDs to experimental human subjects cannot reasonably be performed, some of the data included in this review are from studies of anesthetized animals. Most of the human studies of ECD exposures were supported by

TASER International and involved shorter-duration exposures than some cases of deaths-in-custody reported to be possibly associated with use of the devices.

2. Is “excited delirium” a valid term?

Four components are usually considered to be necessary for a definition of the excited delirium syndrome: a) delirium with agitation, b) respiratory arrest, c) hyperthermia, and d) death.¹ Although some have referred to excited delirium as a “non-medical term,”² it is a valid general description of a syndrome, prior to further determination of an underlying diagnosis.³ Excited delirium can be considered a heterogeneous condition, with multiple variant types that could differ in both presentation and clinical course. The syndrome is properly defined as a physical state for which several underlying diagnoses are feasible,^{4,5,6} with cocaine-induced excited delirium perhaps the most common variant. Although certain clinical cases may be consistent with this sub-type, expert witnesses in some legal cases have suggested that the term “cocaine-induced excited delirium” would not be valid without respiratory arrest and death.⁷ In other instances, some behavioral symptoms may be similar to those observed in psychiatric conditions such as catatonia.⁸ These cases, however, are not

* Tel.: +1 210 536 3572; fax: +1 210 536 3977.

E-mail address: james.jauchem@brooks.af.mil.

necessarily classified, *per se*, as excited delirium. Detweiler et al.⁹ noted that, when a subject exhibits an increasing degree of catatonic features, delirious mania may convert to a malignant catatonic state, particularly in the presence of hyperthermia. (Also see additional discussion in section “7. Body temperature” below.)

As DiMaio and DiMaio¹⁰ pointed out, the fact that excited delirium is not a billable procedure (listed by the American Medical Association) does not mean it is not a valid diagnosis. The authors pointed out that the National Association of Medical Examiners has recognized excited delirium as a valid entity for a number of years. In addition, the American College of Emergency Physicians formally recognized the syndrome recently.¹¹

3. Hyperkalemia

3.1. Hyperkalemia during ECD experiments

As mentioned in the previous review,¹ increased blood potassium concentration ($[K^+]$) was noted in an animal model (*Sus scrofa*) after repeated or long-duration ECD exposures. In subsequent experiments, performed with the same methods of anesthesia and physiological measurements,^{12,13,14} other repeated or long-duration ECD applications also resulted in significant hyperkalemia immediately after exposure. The most extreme hyperkalemia (mean value of 7.6 mEq/L) was seen after 3 min of cycled (7 s on/3 s off) applications of the TASER® X26 ECD.¹² (“TASER” is a registered trademark of TASER International. “X26” is a trademark of TASER International.) Within 30 min, however, potassium had returned to a normal level. There were no significant differences in pre-exposure blood $[K^+]$ from survivors versus non-survivors.

Valentino et al.^{15,16,17} reported no significant change in blood $[K^+]$ for up to 30 days post-exposure in swine due to applications of an Aegis ECD. The same group of investigators,¹⁸ however, found that two 40-s applications of a TASER X26 ECD resulted in a significant increase in blood $[K^+]$ at 5 min post-exposure. In another series of 40-s X26 ECD applications, succinylcholine was administered to swine to prevent acidemia due to muscle contraction.¹⁹ Fatal ventricular fibrillation (VF) occurred in only one animal out of eight exposed to two applications. The authors reported no blood potassium values outside a normal range in their experiments.

In experiments of human volunteers, there were no significant changes in venous blood $[K^+]$ after 5-s applications of the TASER X26 ECD.²⁰ Ho et al.²¹ found no significant changes in serum $[K^+]$ after 15-s TASER X26 “drive stun” applications in human volunteers. These experiments were funded by TASER International. Longer exposure periods were not investigated.

Even though muscle contraction responses to ECD applications would typically bypass volition, the overall responses may be similar to changes during exercise.²² Due to these similarities, knowledge of prior studies of exercise/muscle contraction may be useful in understanding responses during ECD applications. Blood $[K^+]$ will increase to a greater extent in untrained human subjects than in exercise-trained subjects.²³ In addition, after exertion, trained subjects recover to resting concentrations of plasma potassium more quickly.²⁴ Knochel et al.²³ hypothesized that such subjects may be able to avoid potentially dangerous hyperkalemia via biologic adaptation of skeletal muscle cells. Whether or not exercise-trained subjects are better able to tolerate long-duration exposures to ECDs is unknown. Since most muscles would become inactive immediately after an ECD discharge is discontinued, a rapid uptake into the muscle (similar to what may happen after cessation of exercise²⁵) may help prevent any harmful level of extended hyperkalemia.

Hyperkalemia can occur in association with hyperglycemia.²⁶ (See section “8.2. Increased blood glucose after ECD applications or

excited delirium” below, for further discussion relative to ECD applications or excited delirium.) Hyperkalemia is a potentially severe complication of rhabdomyolysis (which may occur after repeated muscle contractions) (also see additional discussion in section “5. Rhabdomyolysis and muscle enzymes” below), and could result in life-threatening arrhythmias.²⁷ In addition to release of potassium by muscle, other factors can play a role in changing blood potassium during muscular exercise. For example, the liver can exchange potassium actively due to activation of adrenergic receptors,²⁸ which could arise due to increased catecholamines.

Although severe hyperkalemia can occur in some patients without major electrocardiographic manifestations,²⁹ profound effects on cardiac conduction (as specifically reflected in the electrocardiogram) usually result.³⁰ Efforts to use quantitative criteria to define effects of hyperkalemia on cardiac conduction have been difficult to validate in humans.³¹ In swine, an increase in T-wave amplitude does not usually happen until blood $[K^+]$ reaches a level of about 7.6 mEq/L.³² In the previous ECD study in which blood $[K^+]$ reached that level,¹² T-wave elevation was not noted. Cardiac arrest will usually take place at a level of 12 mEq/L.³³ Blood $[K^+]$ may not be predictive of hemodynamic instability in any individual patient, since the effects will depend on both the rate of increase and associated metabolic factors, including the pH.³⁴

3.2. Hyperkalemia during excited delirium

Some deaths after cocaine use (without concurrent exposure to ECDs) have been related to severe hyperkalemia (e.g., see Parks et al.³⁴). This aspect may be of interest in selected cases of deaths occurring after excited delirium. In spontaneously breathing pigs (without any ECD applications), administration of cocaine resulted in an increase in blood $[K^+]$ from 3.9 mEq/L pre-infusion to 6.5 mEq/L 2 h after start of infusion.³⁵ The authors, however, suggested that “respiratory depression and resulting hypoxemia and acidosis were the primary reasons for cardiac arrest” in the animals. As in ECD experiments,¹² changes in respiration and acidosis (rather than hyperkalemia) may also have been more important factors relating to death during excited delirium. (Also see additional discussion in section “6. Respiration, acidosis, and blood lactate” below.)

In cases of life-threatening hyperkalemia in humans due to methamphetamine³⁶ or cocaine,³⁷ blood $[K^+]$ levels were 8.5 and 8.9 mEq/L, respectively. In one recent case of rhabdomyolysis due to 3,4-methylenedioxy-N-methylamphetamine (MDMA, “ecstasy”), hyperkalemia (blood $[K^+]$ of 9.4 mEq/L) was noted as the cause of death.³⁸ Since drug abuse is often present in cases of excited delirium (or, additionally, ECD usage), these associations should be considered. (Also see additional discussion in section “9. Additional information regarding drug abuse” below.)

4. Direct and indirect effects of ECDs or excited delirium on the cardiovascular system

4.1. ECDs

Using human subjects in experiments funded by TASER International, Ho et al.³⁹ reported no detectable changes in 12-lead electrocardiograms due to 15-s TASER X26 ECD exposures immediately after an exhaustive exercise regimen. In another study of up to 5-s X26 device applications, the exposures appeared “to be safe and well tolerated from a cardiovascular standpoint.”⁴⁰ Heart rate and mean blood pressure were increased slightly but significantly. Vilke et al.⁴¹ also reported increased heart rate, but no changes in ventilatory parameters, after 5-s X26 ECD exposures. Dawes et al.⁴² found no cardiac capture in ten volunteer subjects exposed to 5-s X26 ECD discharges, with the probes actually fired into the chest

region (and not simply taped to conductive gel, as in previous studies of humans). After 10-s applications of a “new generation” ECD manufactured by TASER International, there were no apparent electrocardiogram changes of any importance.⁴³ (In experiments of an earlier preliminary prototype version of the device,⁴³ one subject demonstrated rapid ventricular capture for the duration of the discharge.) These last two studies mentioned above^{42,43} were funded by TASER International. The findings may not apply to situations of ECD applications of longer durations or with repetitive discharges.

In arrest-related deaths occurring in persons who collapsed less than 15 min after ECD application and in whom rhythm was documented, asystole or pulseless electrical activity was the rhythm primarily found.⁴⁴ The findings of rhythm analysis are a necessarily small group and the sample size is worthy of caution. While none of the 8 subjects who collapsed (while actually on a cardiac monitor) demonstrated VF, one other subject's demise was one of the first recorded cases of VF potentially induced by ECD application.

Investigators have presented convincing evidence that, under certain selected conditions, ECD applications can result in ventricular capture (and associated mechanical contraction) in an anesthetized swine model.^{18,19,45} Whether different anesthetics play a role in such responses is unknown (see Jauchem⁴⁶ for discussion of such potential effects). In addition, there is controversy regarding the appropriateness of a swine model for studying susceptibility to arrhythmias, with some researchers affirming this idea (reviewed by Jauchem^{46–47}) and others disputing it.⁴⁸ Ho et al.⁴⁹ reported that ECD applications resulted in greater atrial and ventricular irritability in methamphetamine-intoxicated Dorset sheep (*Ovis aries Dorset*) of small body weight. Despite these findings, cardiac events such as VF are still unlikely to occur, in most cases, as a result of any direct effect of ECD pulses on the heart.⁵⁰ There was no VF in animals that expired after extreme ECD exposures,¹² but rather respiratory arrest with organized QRS electrical activity. Thus there was possibly pulseless electrical activity. (When such activity does occur, it may change spontaneously to asystole.⁵¹)

Cevik et al.⁵² suggested an ECD “application does not have an acute direct effect on cardiac function but causes a final sympathetic surge from the central nervous system.” Although not based on any data of any specific research studies on the topic, Holden et al.⁵³ speculated that, independently of any primary electrical effect of ECDs, arrhythmias may be caused by exercise- or stress-induced catecholamine release. Hyperkalemia, in combination with high catecholamine levels, could actually facilitate cessation of VF and, therefore, be protective.⁵⁴ (Also see additional discussion in section “10. The potential combination of factors” below.)

Michaud and Dupuis⁵⁵ hypothesized that “the sudden tetanic muscular contractions” induced by ECDs “represent a high level of mechanical energy that is nothing less than a severe blunt trauma,” which could result in dysrhythmias due to “commotio cordis.” (Coincidentally, a swine model is appropriate for studying such a cause of dysrhythmia.⁵⁶)

In experiments of 30- or 60-s ECD applications in pigs,¹³ detectable levels of cardiac troponin I were present in only one out of ten animals. Ho et al.²¹ reported that all types of troponins measured were <0.2 ng/ml immediately, 8 h, and 24 h after TASER X26 “drive stun” applications in human volunteers. VanMeenen et al.⁵⁷ also found no detectable levels of cardiac troponin I, in a group of 99 law-enforcement officers and trainees, 24 h after 5-s X26 ECD exposures.

4.2. Excited delirium

Roberts⁵⁸ noted “previous reports have documented PEA/asystole rather than ventricular tachycardia or fibrillation” in “patients with cocaine-induced delirium who suffered sudden unexpected cardiac arrest.” Since cocaine is often present in subjects exposed to

ECDs during law-enforcement incidents,¹ this is an important observation. Long-term cocaine use, however, is necessary for the development of cardiac myofibrillary alterations.⁵⁹ In persons aged 18–45, more than ten uses of cocaine significantly increases the risk of myocardial infarction, particularly in younger persons.⁶⁰ While myofibrillary changes may not be seen on pathology until after long-term cocaine use, the clinical effect of cocaine and coronary insufficiency are seen early in young and seemingly healthy people with no other risk factors.⁶¹ Concomitant use of cigarettes and alcohol potentiates the cardiac risks of cocaine.⁶²

5. Rhabdomyolysis and muscle enzymes

5.1. ECDs

In experiments of 30- or 60-s ECD applications in pigs,¹³ serum myoglobin was increased significantly immediately after exposure and at all sampling points in a 3-h post-exposure period. In another series of cycled applications (7 s on/3 s off) of the X26 ECD,¹² serum myoglobin was also increased significantly immediately after exposure and at most sampling points post-exposure. The maximum value (76 ng/ml), however, was less than half that which occurred in a study of healthy men performing isometric or isokinetic exercise.⁶³

Sanford et al.⁶⁴ presented two cases of men with rhabdomyolysis after exhibiting violent behavior and receiving TASER X26 ECD applications. One of the subjects had received six ECD applications and had a serum myoglobin of 169 ng/ml. The other subject (after only a single ECD shock) exhibited a peak creatine phosphokinase (CPK) of 8086 units. As Ho and Dawes⁶⁵ pointed out, however, the relative contributions of physical struggle versus ECD applications in cases of this nature have not been determined. Other investigators have found that exhaustive muscle contractions (e.g., 50 maximal contractions of elbow flexor muscles⁶⁶) result in large increases in blood CPK.

Significant increases in serum total CPK and in the CPK-MM fraction have occurred after 30- or 60-s ECD exposures,¹³ and even after only three 5-s applications of a TASER X26 ECD,⁶⁷ to anesthetized swine. Sloane and Vilke⁶⁸ suggested that increases in CPK would not be surprising after ECD application, since massive contraction of muscle could cause such changes. On the basis of pooled data from several different studies of humans (a meta-analysis funded by TASER International), Dawes et al.⁶⁹ reported no clinically significant changes in CPK after ECD exposures (in contrast with the X26 ECD results in animal studies). The vast majority of the human studies, however, involved only a single 5-s application. It is unknown whether longer applications to humans would result in clinically significant rhabdomyolysis.

5.2. Excited delirium

Although verified cases of strictly-defined excited delirium in conjunction with rhabdomyolysis were not studied, the most recent report of rhabdomyolysis due to licit or illicit drugs was by Tóth and Varga.⁷⁰ Drugs included amphetamine, cocaine, alcohol, MDMA, heroin, and methadone. Berrens et al.⁷¹ presented a case of rhabdomyolysis caused by D-lysergic acid diethylamide.

6. Respiration, acidosis, and blood lactate

6.1. ECDs

Respiration is dependent on a “bellows” mechanism via the action of the abdominal wall, diaphragm, intercostal muscles, and accessory muscles of respiration.⁷² During ECD applications, each of

these could be affected. Gill⁷³ suggested that prolonged ECD applications could cause respiratory problems. Studies of ECD effects on respiration were reviewed previously.⁴⁷ A lack of effective respiration, which was seen in anesthetized animals, has not been replicated in experiments of human subjects. Dawes et al.,⁷⁴ however, noticed in prior studies “that subjects often held their breath for the first 5–7 s due to a pain response” to ECD exposure. Human studies are limited by the need to include healthy individuals in clinically-controlled research settings. In some studies, the subjects may have been aware of the need to actively focus on breathing.⁷⁵

Vilke et al.²⁰ found a doubling of arterial blood lactate concentrations in human subjects exposed to a single 5-s application of the X26 ECD. Other similar studies have been reviewed previously.⁴⁷ In a study of ethanol-intoxicated subjects,⁷⁶ a 15-s TASER X26 ECD discharge resulted in a greater rate of increased blood lactate (and decreased pH) than during the pre-exposure period. Ho et al.⁷⁷ reported that a 15-s exposure to the X26 ECD did not worsen acidemia that was already present in anaerobically exhausted human subjects, although there was a slight increase in blood lactate. In another study,⁷⁸ a similar ECD exposure did not appear to worsen acidemia in exhausted subjects any differently than continued exertion. (These last three studies mentioned above were funded by TASER International.) The investigators noted that their data may not apply to situations of ECD applications of longer durations or repetitive discharges. Lactic acidosis resulted from exposures to an ECD marketed for civilians (“TASER C2”) in experiments of both human⁷⁹ and animal¹⁴ subjects. Although some investigators may consider acidosis to be “a life-threatening condition regardless of the underlying condition,”⁸⁰ short ECD applications may cause only very transient decreases in blood pH.

6.2. Excited delirium

Whether asphyxia is associated with excited delirium is uncertain. Byard et al.⁸¹ have reviewed studies pertinent to this question. Hypoxia, even in anesthetized animals (i.e., without the perception of pain), could trigger a rapid stress reaction, leading to the release of both epinephrine and norepinephrine into the blood. Some postmortem findings associated with asphyxia are nonspecific.⁷² In cases of asphyxiation, blood levels of epinephrine and norepinephrine can increase immediately at death.⁸² Hirvonen et al.⁸³ determined that increased postmortem blood concentrations of catecholamines could be used as indicators of hypoxia, with the caveat that levels could rise in a short period after death. Takeichi et al.⁸² reported that epinephrine may gradually decrease, while norepinephrine may increase further, due to release postmortem from cardiac sympathetic nerves. All of these results may be heavily confounded by other events that increase catecholamine levels independently from asphyxia, i.e., stress, agitation, exercise, acidosis, and a host of other issues that could be concomitant. Acidosis can directly stimulate release of epinephrine from the adrenal medulla.⁸⁴ (Also see section “10. The potential combination of factors” below.)

7. Body temperature

No significant increases in body temperature were found after extreme applications of the TASER X26 device¹² or after 30-s applications of the TASER C2 ECD¹⁴ using an animal model. Lim and Seet⁸⁵ were surprised, considering the vulnerability of nervous tissue to thermal injury, that neurological effects were not reported more frequently after ECD exposures. Although Bui et al.⁸⁶ recently reported neurological sequelae of ECD applications, these were not thermal in nature. In a population-based study of 426 ECD

activations by law-enforcement personnel,⁸⁷ only one death was subsequently attributed to hyperthermia.

Hyperthermia is a pivotal issue in discussions surrounding excited delirium, since the presence of hyperthermia in a state of excited delirium is considered a harbinger of death.⁸⁸ As Bunai et al.⁸⁹ noted, “Although the findings associated with hyperthermia” during excited delirium “are nonspecific, they are useful in diagnosing hyperthermia when combined with reconstruction of the scene of death.” The most recent report of a case of cocaine-induced excited delirium, concurrent with hyperthermia, was by Menaker et al.⁹⁰ An increase in body temperature at time of death could be caused by a variety of factors, including physical activity, emotional stress, intoxication due to illicit drugs, and hyperthyroidism.⁹¹ Each of these may be relevant to excited delirium. (Also see section “11.5 Hyperthermia” below.)

8. Additional topics not discussed in previous review: Increases in hematocrit and blood glucose

8.1. Increased hematocrit after ECD applications or excited delirium

In one ECD study of swine, non-survivors exhibited a significantly higher pre-exposure hematocrit than survivors.¹² In all previous^{67,92} and subsequent^{13,14} ECD experiments in which hematocrit was measured, the value also increased significantly after exposure. The mechanisms of these changes in hematocrit have not been determined, but potential candidates have been reviewed previously.⁹³ These include: a) increased capillary hydrostatic pressure due to effects of muscle contraction on venules (with plasma fluid pushed into the extravascular space); b) release of metabolic waste products from muscles (with increased osmolality of the interstitial space); and c) splenic contraction.

Potential detrimental effects of acutely-increased hematocrit (and related hemorheological alterations⁹⁴) include problems with tissue perfusion, and reduced venous return and cardiac output due to higher viscosity.⁹³ Although polycythemia increases blood oxygen content, this potential benefit may be counterbalanced by a decrease in cardiac output, which will, in turn, adversely affect pulmonary diffusion and alveolar ventilation.

In a study of 53 human subjects exposed for 10 s to a particular ECD (the same model as mentioned in another reference⁴³ above), there was no change in mean hematocrit.⁹⁵ Changes of hematocrit in humans exposed to long-duration or repeated applications of ECDs, however, have not been studied. Unfortunately, since hematocrit values increase rapidly after death,⁹⁶ a forensic investigation of this phenomenon would be problematic. On the basis of the information presented above, however, the clinical implications of increased hematocrit could be serious. In addition, although there have been no published reports of autopsy findings of hypercoagulability or clot-related demise after ECD applications, acute muscle contraction (even without increased hematocrit) can result in increased thrombotic tendency,⁹⁷ with sustained platelet and coagulation hyperactivity.⁹⁸ Acute muscle contraction can result in both prothrombic and hypofibrinolytic responses.⁹⁹ Any such detrimental effects, however, would appear intuitively to be dependent on a high level of either long-duration or repetitive ECD applications. Low degrees of muscle contraction may not have such effects.

Cocaine can cause a transient increase in hematocrit in humans, due to splenic contraction.¹⁰⁰ If a subject has pre-existing polycythemia from this or other causes (e.g., an endocrinopathy¹⁰¹), ECD application could exacerbate the situation. Although, in rare cases, fatalities that include cocaine-associated rhabdomyolysis may be associated with disseminated intravascular coagulation,¹⁰² such a process would likely occur only after some time period,

e.g., during the occurrence of renal failure.¹⁰³ Thus it is unlikely that ECD applications would be related to any cases of disseminated intravascular coagulation.

8.2. Increased blood glucose after ECD applications or excited delirium

In previous experiments,^{12–14,67,92} significant increases in blood glucose were observed either immediately after or within 30 min after ECD applications. In the study of the most extreme ECD applications,¹² blood glucose increased significantly in non-survivors, but not in animals that survived. These results were consistent with Asadollahi et al.,¹⁰⁴ who noted significantly higher mortality in critically-ill human patients with plasma glucose levels greater than an upper cut-off value of 125 mg/dL (which was identical to the mean value of blood in animals that survived ECD applications in the previous study by Jauchem et al.¹²). In a more recent study, polytraumatized human patients exhibited greater mortality with higher blood levels of glucose¹⁰⁵ compared with normoglycemic individuals. It should be noted, however, that other factors during critical illness and trauma may have no bearing on situations involving deaths-in-custody.

In some acute pathophysiological states, combinations of several different blood factors may be important. For example, hyperglycemia and hyponatremia together may have a cumulative effect on mortality.¹⁰⁴ In experiments with significant increases in blood glucose after ECD application,^{12,13,14,67,92} however, hyponatremia occurred in only one study.⁶⁷ Hyperglycemia may result in decreased erythrocyte deformability,¹⁰⁶ which can occur concurrently with increased hematocrit.⁹³

It is unknown whether increased blood glucose is a function of physical struggle, excited delirium, drug intoxication, or ECD applications during law-enforcement situations. Alcohol,¹⁰⁷ opioids, and opioid analogs¹⁰⁸ can, by themselves, result in hyperglycemia. (Also see additional discussion in section “11.3 Hyperglycemia” below.)

9. Additional information regarding drug abuse

Although deaths due to cocaine occur only in a minority of subjects who use the substance, it is toxic over a large range of blood concentrations.¹⁰⁹ Elevated catecholamines could increase the toxicity of cocaine.¹¹⁰ Pilgrim et al.¹¹¹ confirmed earlier results of Karch et al.¹¹² of no correlation between blood methamphetamine and outcome. The authors¹¹¹ suggested that femoral blood be taken so that artifacts due to postmortem redistribution could be diminished. They noted that amphetamine-class drugs either caused or exacerbated pre-existing cardiovascular disease. Karch⁵⁹ explained that pathological changes in the heart (sometimes seen after excited delirium) due to methamphetamine or cocaine usually require chronic use of the drugs over long periods of time.

Each of the potential effects of factors discussed in the previous sections can be exacerbated by recreational use of cocaine, methamphetamine, or alcohol. Sudden deaths while in custody are nearly always associated with the presence of positive drug levels.

10. The potential combination of factors

Predicting the effects of any given ECD exposure or excited delirium scenario (alone or in combination with each other) is complicated. The combination of acidosis, hyperkalemia, and elevated adrenergic activity may result in antagonism that offsets any harmful effects of an individual factor.¹¹³ O'Neill and Pateron¹¹⁴ suggested that, although hyperkalemia and increased circulating catecholamines individually may each be harmful, there

could actually be a *beneficial* interaction between these changes if they are concurrent. For example, during exercise, catecholamines may modulate calcium currents in the heart, and thereby help to protect it from hyperkalemia.¹¹⁵

Deaths during law-enforcement custody have been likened to a “perfect storm” of severe physiologic conditions including a hyperactive stress response potentiated by restraint, physical exertion, or drug use.¹¹⁶ In this situation, prior health problems may be unmasked. (It is unknown, however, whether ECD exposure would exacerbate any pre-existing conditions.) For example, polycythemia (or associated changes) (also see section “8.1. Increased hematocrit after ECD applications or excited delirium” above) could be present during apprehension of unlawful subjects. Potential causes (prior to any use of an ECD) include psychological stress, endocrinopathy, diabetes, chronic alcohol abuse, cocaine use, or use of opiates.⁹³ There is also an increased lethality of cocaine when used in combination with heroin, opiates, and alcohol.¹⁰⁹

11. Possibilities for future forensic investigations

11.1. General concepts

If some of the pathologic changes outlined above have not been specifically investigated, they each cannot be assumed to have *not* been present. Even if such changes are found, they may be taken out of context when considering the whole situation. There are few solutions for elucidating causes of deaths-in-custody. Some of the points below, however, may provide ideas for future investigations.

11.2. Blood catecholamines

Blood should be drawn from the subclavian or femoral vessels to prevent postmortem redistribution of drugs from tissues into blood, which could result in elevated levels as artifacts.¹¹⁷ If a relatively long period of time has elapsed, blood obtained at the beginning of admission to a hospital (if available) could still be valuable for toxicologic analysis.¹¹⁸

High levels of catecholamines in the circulation could lead to hyperthermia and increased metabolism. If these changes are not apparent on autopsy, a lack of findings could occur in some cases of deaths-in-custody.¹¹⁹ Often, no blood samples are available immediately after cases of excited delirium.

Postmortem levels of blood catecholamines may be related to the magnitude of stress responses in individuals.¹²⁰ Elevated levels of catecholamines postmortem in vitreous humor and cerebrospinal fluid could be associated with a long-lasting stress reaction,¹²¹ such as that occurring during excited delirium.

11.3. Hyperglycemia

Epinephrine simultaneously enhances hepatic glucose production and inhibits glucose clearance.¹²² Increased sympathetic drive during ECD exposures or excited delirium may be important in causing increases in blood glucose. Zilg et al.¹²³ suggested that, since blood glucose levels decrease rapidly postmortem, vitreous humor glucose alone should be used to determine hyperglycemia. Possible postmortem changes such as redistribution and hemolysis, however, should be considered.¹²⁴

11.4. Inherited ion channelopathies

After some deaths-in-custody, postmortem analysis for inherited ion channelopathies¹²⁵ could be considered. With recent advances in molecular technology, examination of cardiac-channel

factors may provide a potential pathogenic basis for unexplained deaths.^{126–127} Examination of myocardial contraction band necrosis potentially due to acute use of drugs (e.g., MDMA),^{128–129} would also be of particular interest.

11.5. Hyperthermia

Since hyperthermia is often evident during excited delirium (versus a lack of increased temperature during ECD exposures), a major focus for new forensic investigations may involve this factor. By examining dopamine dysregulation biomarkers in the brain postmortem, and by analyzing induction of heat shock protein, Mash et al.⁸⁸ found that death likely occurs in conjunction with hyperthermia during excited delirium. Concentrations of dopamine transporters have been below a normal range in all excited delirium cases in which such levels have been examined.¹³⁰ Isolated increases in serum creatinine levels may be useful in investigations of death due to hyperthermia,¹³¹ which often includes absent or nonspecific findings at autopsy.¹³² In addition to these factors, other aspects or markers that may distinguish excited delirium hyperthermic effects from ECD effects (i.e., events that occur with hyperthermia but probably *not* with ECD applications alone) include: elevated cardiac troponin T in left and right heart blood, external iliac blood, pericardial fluid,¹³³ and cerebrospinal fluid¹³⁴; elevated urea nitrogen, creatinine, and uric acid in pericardial fluid¹³⁵; elevated calcium (but decreased magnesium) in pericardial fluid¹³⁶; decreased adrenocorticotrophic hormone in cerebrospinal fluid¹³⁷; increased ubiquitin immunoreactivity in midbrain periaqueductal gray matter¹³⁸; elevated mast cell tryptase in serum¹³⁹; increased ratio of serum/cerebrospinal chromogranin A¹⁴⁰; and lower neuronal dopamine-immunopositivity in the hypothalamus.¹⁴¹

12. Clinical context

Since complete statistics relating to all deaths-in-custody are not available in the public domain,¹⁴² research in this area may be problematic. Hall¹⁴³ emphasized the importance of placing potential risks of ECD usage during law-enforcement activities into proper perspective with other alternatives. She suggested that collection of data related to such use should take precedence over studies of animal models or healthy human subjects.

Notwithstanding results of animal experiments with direct effects on the heart in some specific conditions of TASER ECD applications (e.g., Valentino et al.⁴⁵), the likelihood of ventricular VF *directly* due to such devices has been reported to be extremely low during typical fatality cases involving excited delirium.¹⁴⁴ In one study of deaths-in-custody between 1990 and 2004, only one out of 45 cases involved use of an ECD.¹⁴⁵ In a prospective multicenter cohort study of 1201 subjects exposed to ECDs, two deaths-in-custody occurred; both were determined to be unrelated to ECD use.¹⁴⁶ In contrast, on the basis of some analyses, as many as 75% of excited delirium patients die at scene or during transport (C. Hall, quoted in Calgary Police Commission¹⁴⁷). Bozeman and Winslow¹⁴⁸ concluded that drugs and/or restraint may be more important factors leading to death than several short exposures to ECDs. In 2006, it was estimated that as many as 800 people die from excited delirium each year.¹⁴⁹ Since different terminology and definitions of excited delirium may be used by different medical examiners, there are no reliable figures nationwide of suspects who died from the syndrome. One investigator estimated an in-custody death occurrence approximately every 2.5 days in the United States, with ECDs used on these subjects about 27% of the time.¹⁵⁰

The very group of potential subjects exposed to ECDs that may be the most important, in terms of susceptibility to death (i.e., those

exhibiting excited delirium), cannot be studied ethically in controlled experiments. Studies of volunteers might not be directly extrapolatable to law-enforcement situations with excited delirium or restraint of subjects.¹⁵¹ This factor is a challenge in terms of potential research project design. Whether more extensive videotaping, etc., during law-enforcement use of ECDs will greatly improve our understanding is unknown.

Grant et al.¹⁵² noted, “relevant data on the length of time an individual was in custody before death occurred were not consistently available.” The length of time from ECD applications until death has varied from minutes to days. Gill⁷³ remarked that ECDs “certainly cause pain and stress.” It could be reasonable, regarding individuals with advanced heart disease, to consider these external stressors, along with the time sequence of events and other circumstances. It may not be possible, however, to either prove or refute applications of an ECD as a direct cause of death in a subject exhibiting excited delirium.⁷²

Relationships between excited delirium and other events occurring during deaths-in-custody (e.g., manual restraint and oleoresin capsicum) are beyond the scope of this paper, but have been reviewed previously by other investigators.^{153–154} (Use of a new term, “cardiac dysrhythmia during restraint,” was recently suggested as “the most intellectually honest cause of death in some challenging cases” of deaths-in-custody.¹⁵⁵)

As noted previously, “it is possible that there are no unique features of TASER ECD applications, in terms of physiological effects, other than the extreme degrees of muscle contraction (either repeated or long-duration) that may be achieved.”²²

13. Conclusions

Differences between changes in factors due to excited delirium versus those of ECD applications are not readily apparent in most cases. These factors include: direct and indirect effects of the cardiovascular system, respiration, rhabdomyolysis and muscle enzymes, hyperkalemia, acidosis, hyperglycemia, and increased hematocrit. On the basis of this review, however, more complete examination of pathological changes due to hyperthermia (which is present during excited delirium, but not during ECD exposures) may be useful to distinguish between the two different entities. Despite the potential value of this approach, deaths-in-custody may not be solely due to either of these events by themselves, but must always be considered in the context of all possible interacting factors.

Conflict of interest

The author has not had any relationship with any manufacturers of electronic control devices, including employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding.

Acknowledgments

The views, opinions and/or findings contained in this report are those of the author and should not be construed as an official Department of Defense position, policy or decision.

References

1. Jauchem JR. Deaths in custody: are some due to electronic control devices (including TASER® devices) or excited delirium? *J Forensic Leg Med* 2010;17:1–7 [review].
2. Pearson SS. TASER use is ‘organized bullying.’ *Emerg Med News* 2008;30(4):3 [letter].
3. Hall C. Excited delirium: a real medical condition. *Can Police Assoc Express* 2008;71:14–5 (Winter).

4. Hall CA. Sudden death proximal to police restraint: medical issues. Canadian Association for Civilian Oversight of Law Enforcement. Conference. Montreal, Quebec, Canada: Current Issues in Police Accountability; October 3–5, 2005.
5. Hall C, Butler C. National study on neck restraints in policing. Technical report TR-03-2007. Ottawa, Ontario, Canada: Canadian Police Research Centre; June 2007.
6. Stuart B, Lawrence C, Hall C. Report on conducted energy weapons and excited delirium syndrome. Ottawa, Ontario, Canada: Royal Canadian Mounted Police; October 29, 2007.
7. Leong GB, Leisenring SE, Dean MD. Commentary: intoxication and settled insanity — unsettled matters. *J Am Acad Psychiatry Law* 2007;**35**:183–7.
8. Samuel E, Williams RB, Ferrell RB. Excited delirium: consideration of selected medical and psychiatric issues. *Neuropsychiatr Dis Treat* 2009;**5**:61–6.
9. Detweiler MB, Mehra A, Rowell T, Kim KY, Bader G. Delirious mania and malignant catatonia: a report of 3 cases and review. *Psychiatr Q* 2009;**80**:23–40.
10. DiMaio VJM, DiMaio TG. Excited delirium syndrome. In: Kroll MW, Ho JD, editors. *TASER® conducted electrical weapons: physiology, pathology, and law*. New York: Springer Science+Business Media; 2009. p. 347–63.
11. Hoffman L. ACEP recognizes excited delirium syndrome. *Emerg Med News* 2009;**31**(10) (October 5).
12. Jauchem JR, Seaman RL, Fines DA. Survival of anesthetized Sus scrofa after cycling (7 s on / 3 s off) exposures to a TASER® X26 electronic control device for three minutes. *Am J Forensic Med Pathol*; in press.
13. Jauchem JR, Beason CW, Cook MC. Acute effects of an alternative electronic-control-device waveform in swine. *Forensic Sci Med Pathol* 2009;**5**:2–10.
14. Jauchem JR, Seaman RL, Klages CM. Physiological effects of the TASER® C2 conducted energy weapon. *Forensic Sci Med Pathol* 2009;**5**:189–98.
15. Valentino DJ, Walter RJ, Nagy K, Dennis AJ, Winners J, Bokhari F, et al. Repeated thoracic discharges from a stun device. *J Trauma* 2007;**62**:1134–42.
16. Valentino DJ, Walter RJ, Dennis AJ, Nagy K, Loo MM, Winners J, et al. Neuromuscular effects of stun device discharges. *J Surg Res* 2007;**143**:78–87.
17. Valentino DJ, Walter RJ, Dennis AJ, Nagy K, Loo MM, Winners J, et al. Acute effects of MK63 stun device discharges in miniature swine. *Milit Med* 2008;**173**:167–73.
18. Dennis AJ, Valentino DJ, Walter RJ, Nagy KK, Winners J, Bokhari F, et al. Acute effects of TASER X26 discharges in a swine model. *J Trauma* 2007;**63**:581–90.
19. Walter RJ, Dennis AJ, Valentino DJ, Margeta B, Nagy KK, Bokhari F, et al. TASER X26 discharges in swine produce potentially fatal ventricular arrhythmias. *Acad Emerg Med* 2008;**15**:66–73.
20. Vilke GM, Sloane CM, Bouton KD, Kolkhorst FW, Levine SD, Neuman TS, et al. Physiological effects of a conducted electrical weapon on human subjects. *Ann Emerg Med* 2007;**50**:569–75.
21. Ho JD, Dawes DM, Lapine AL, Heegaard WG, Bultman LL, Johnson MA, et al. Prolonged TASER® “drive stun” exposure in humans does not cause worrisome biomarker changes. Abstracts for the 2008 National Association of EMS Physicians (NAEMSP) Scientific Assembly. *Prehosp Emerg Care* 2008;**12**:128 [abstract].
22. Jauchem JR. Muscle stimulation by TASER® conducted energy weapons: similarities with voluntary muscle contractions during exercise. *Adv Med Biol* 2010;**7**:169–79.
23. Knochel JP, Blachley JD, Johnson JH, Carter NW. Muscle cell electrical hyperpolarization and reduced exercise hyperkalemia in physically conditioned dogs. *J Clin Invest* 1985;**75**:740–5.
24. Marcos E, Ribas J. Kinetics of plasma potassium concentrations during exhausting exercise in trained and untrained men. *Eur J Appl Physiol Occup Physiol* 1995;**71**:207–14.
25. Medbø JJ, Sejersted OM. Plasma potassium changes with high intensity exercise. *J Physiol* 1990;**421**:105–22.
26. Mahoney BA, Smith WA, Lo DS, Tsoi K, Tonelli M, Clase CM. Emergency interventions for hyperkalemia. *Cochrane Database Syst Rev* 2005;(2). CD003235 [review].
27. Mohaupt MG. Rhabdomyolysis. *Ther Umsch* 2003;**60**:391–7 [German].
28. Jauchem JR, Vick RL. Phenylephrine-induced hyperkalemia: role of the liver. *Proc Soc Exp Biol Med* 1980;**163**:478–81.
29. Martinez-Vea A, Bardaji A, Garcia C, Oliver JA. Severe hyperkalemia with minimal electrocardiographic manifestations: a report of seven cases. *J Electrocardiol* 1999;**32**:45–9.
30. Diercks DB, Shumak GM, Harrigan RA, Brady WJ, Chan TC. Electrocardiographic manifestations: electrolyte abnormalities. *J Emerg Med* 2004;**27**:153–60 [review].
31. Freeman K, Feldman JA, Mitchell P, Donovan J, Dyer KS, Eliseo L, et al. Effects of presentation and electrocardiogram on time to treatment of hyperkalemia. *Acad Emerg Med* 2008;**15**:239–49.
32. Coulter DB, Swenson MJ. Effects of potassium intoxication on porcine electrocardiograms. *J Vet Res* 1970;**31**:2001–11.
33. Geddes LA, Roeder RA, Rundell AE, Otlewski MP, Kemeny AE, Lottes AE. The natural biochemical changes during ventricular fibrillation with cardiopulmonary resuscitation and the onset of postdefibrillation pulseless electrical activity. *Am J Emerg Med* 2006;**24**:577–81.
34. Parks JM, Reed G, Knochel JP. Cocaine-associated rhabdomyolysis. *Am J Med Sci* 1989;**297**:334–6.
35. Kambam J, Mets B, Hickman R, Janicki P, James MF, Kirsch R. The effects of inhibition of plasma cholinesterase activity on systemic toxicity and blood catecholamine levels from cocaine infusion in pigs. *J Lab Clin Med* 1993;**122**:188–96.
36. Chan P, Chen JH, Lee MH, Deng JF. Fatal and nonfatal methamphetamine intoxication in the intensive care unit. *J Toxicol Clin Toxicol* 1994;**32**:147–55.
37. Siddiqui F, Slater R, Ashraf S. Life threatening hyperkalemia following cocaine ingestion: a case report. *Cases J* 2009;**2**:7355.
38. Eede HV, Montenij LJ, Touw DJ, Norris EM, Rhabdomyolysis in MDMA intoxication: a rapid and underestimated killer. “Clean” ecstasy: a safe party drug? *J Emerg Med*; in press. doi:10.1016/j.jemermed.2009.04.057.
39. Ho JD, Dawes DM, Heegaard WG, Calkins HG, Moscati RM, Miner JR. Absence of electrocardiographic change after prolonged application of a conducted electrical weapon in physically exhausted adults. *J Emerg Med*; in press. doi:10.1016/j.jemermed.2009.03.023.
40. Bozeman WP, Barnes Jr DG, Winslow 3rd JE, Johnson 3rd JC, Phillips CH, Alson R. Immediate cardiovascular effects of the TASER X26 conducted electrical weapon. *Emerg Med J* 2009;**26**:567–70.
41. Vilke GM, Sloane CM, Suffecool A, Kolkhorst FW, Neuman TS, Castillo EM, et al. Physiologic effects of the TASER after exercise. *Acad Emerg Med* 2009;**16**:704–10.
42. Dawes DM, Ho JD, Reardon RF, Miner JR. Echocardiographic evaluation of TASER X26 probe deployment into the chests of human volunteers. *Am J Emerg Med* 2010;**28**:49–55.
43. Ho JD, Dawes DM, Reardon RF, Strode SR, Kunz SN, Nelson RS, et al. Human cardiovascular effects of a new generation conducted electrical weapon. *Forensic Sci Int* 2011;**204**:50–7.
44. Swerdlow CD, Fishbein MC, Chaman L, Lakkireddy DR, Tchou P. Presenting rhythm in sudden deaths temporally proximate to discharge of TASER conducted electrical weapons. *Acad Emerg Med* 2009;**16**:1–13.
45. Valentino DJ, Walter RJ, Dennis AJ, Margeta B, Starr F, Nagy KK, et al. Taser X26 discharges in swine: ventricular rhythm capture is dependent on discharge vector. *J Trauma* 2008;**65**:1478–85.
46. Jauchem JR. An animal model to investigate effectiveness and safety of electronic control devices (including TASER® devices). *J Forensic Sci* 2010;**55**:521–6.
47. Jauchem JR. Repeated or long-duration TASER® electronic control device exposures: acidemia and lack of respiration. *Forensic Sci Med Pathol* 2010;**6**:46–53 [review].
48. Richardson ES. Intrapericardial delivery of anti-arrhythmic agents. PhD Dissertation. University Of Minnesota; May 2009.
49. Ho J, Dawes D, Cole J, Reardon RF, Lundin EJ, Terwey KS, et al. Effect of an electronic control device exposure on a methamphetamine intoxicated animal model. *Acad Emerg Med* 2010;**17**:436–43.
50. Beason CW, Jauchem JR, Clark III CD, Parker JE, Fines DA. Pulse variations of a conducted energy weapon (similar to the TASER® X26 device): effects on muscle contraction and threshold for ventricular fibrillation. *J Forensic Sci* 2009;**54**:1113–8.
51. Skogvoll E, Eftestøl T, Gundersen K, Kvaløy JT, Kramer-Johansen J, Olasveengen TM, et al. Dynamics and state transitions during resuscitation in out-of-hospital cardiac arrest. *Resuscitation* 2008;**78**:30–7.
52. Cevik C, Otahbachi M, Miller E, Bagdure S, Nugent KM. Acute stress cardiomyopathy and deaths associated with electronic weapons. *Int J Cardiol* 2009;**132**:312–7.
53. Holden SJ, Sheridan RD, Coffey TJ, Scaramuzza RA, Diamantopoulos P. Electromagnetic modelling of current flow in the heart from TASER devices and the risk of cardiac dysrhythmias. *Phys Med Biol* 2007;**52**:7193–209.
54. Nair K, Umapathy K, Downar E, Nanthakumar K. Aborted sudden death from sustained ventricular fibrillation. *Heart Rhythm* 2008;**5**:1198–200.
55. Michaud A, Dupuis JY. Echocardiographic evaluation of TASER X26 in healthy volunteers. *Am J Emerg Med* 2010;**28**:521–3 [letter].
56. Weinstock J, Maron BJ, Song C, Mane PP, Estes 3rd NA, Link MS. Failure of commercially available chest wall protectors to prevent sudden cardiac death induced by chest wall blows in an experimental model of commotio cordis. *Pediatrics* 2006;**117**:e656–62.
57. VanMeenen KM, Cherniack NS, Bergen MT, Gleason LA, Teichman R, Servatius RJ. Cardiovascular evaluation of electronic control device exposure in law enforcement trainees: a multisite study. *J Occup Environ Med* 2010;**52**:197–201.
58. Roberts JR. Acute agitated delirium from cocaine: a medical emergency. *Emerg Med News* 2007;**29**(10):18–20.
59. Karch SB. Stimulant abuse and sudden cardiac death. In: Kroll MW, Ho JD, editors. *TASER® conducted electrical weapons: physiology, pathology, and law*. New York: Springer Science+Business Media; 2009. p. 315–25.
60. Aslibekyan S, Levitan EB, Mittleman MA. Prevalent cocaine use and myocardial infarction. *Am J Cardiol* 2008;**102**:966–9.
61. Vasica G, Tennant CC. Cocaine use and cardiovascular complications. *Med J Aust* 2002;**177**:260–2 [review].
62. Afonso L, Mohammad T, Thatai D. Crack whips the heart: a review of the cardiovascular toxicity of cocaine. *Am J Cardiol* 2007;**100**:1040–3.
63. Roxin LE, Venge P, Friman G. Variations in serum myoglobin after a 2-min isokinetic exercise test and the effects of training. *Eur J Appl Physiol Occup Physiol* 1984;**53**:43–7.
64. Sanford JM, Jacobs GJ, Roe EJ, Terndrup TE. Two patients subdued with a TASER® device: cases and review of complications. *J Emerg Med* 2011;**40**:28–32.
65. Ho JD, Dawes DM. TASER device-induced rhabdomyolysis is unlikely. *J Emerg Med* 2011;**40**:68–9.

66. Yamin C, Oliveira J, Meckel Y, Eynon N, Sagiv M, Ayalon M, et al. CK-MM gene polymorphism does not influence the blood CK activity levels after exhaustive eccentric exercise. *Int J Sports Med* 2010;**31**:213–7.
67. Jauchem JR, Cook MC, Beason CW. Blood factors of *Sus scrofa* following a series of three TASER® electronic control device exposures. *Forensic Sci Int* 2008;**175**:166–70.
68. Sloane C, Vilke GM. Riot control agents, tasers, and other less lethal weapons. In: Ross DL, Chan TC, editors. *Forensic science and medicine: sudden deaths in custody*. Totowa, New Jersey: Humana Press, Inc; 2006. p. 113–38.
69. Dawes DM, Ho JD, Miner JR. A meta-analysis of the effect of an electronic control device on muscle injury as determined by creatine kinase enzyme. *Australasian Coll for Emerg Med Scientific Assembly*. Poster available at <http://www.ipcd.com/docs/Med%2520Poster%2520CK.pdf>; November 15, 2009.
70. Tóth AR, Varga T. Myocardium and striated muscle damage caused by licit or illicit drugs. *Leg Med (Tokyo)* 2009;**11**(Suppl 1):S484–7.
71. Berrens Z, Lammers J, White C. Rhabdomyolysis after LSD ingestion. *Psychosomatics* 2010;**51**:356–3563. e3.
72. Shkrum MJ, Ramsay DA. *Asphyxia. Chapter 3. Forensic science and medicine: forensic pathology of trauma: common problems for the pathologist*. Totowa, New Jersey, USA: Humana Press Inc; 2007. pp. 65–179.
73. Gill JR. The medicolegal evaluation of excited delirium. Chapter 6. In: Tsokos M, editor. *Forensic pathology Reviews. Volume 5*. Totowa, New Jersey, USA: Humana Press Inc; 2008. p. 91–111.
74. Dawes DM, Ho JD, Reardon RF, Sweeney JD, Miner JR. The physiologic effects of multiple simultaneous electronic control device discharges. *West J Emerg Med* 2010;**11**:49–56.
75. Reilly JP. *Testimony at the Thomas R. Braidwood, Q.C., Commissions of Inquiry under the public Inquiry Act*. SBC 2007, c. 9. British Columbia, Canada: Vancouver. Available at, <http://www.braidwoodinquiry.ca/transcripts/08-05-05.pdf>; 2008.
76. Moscati R, Ho JD, Dawes DM, Miner JR. Physiologic effects of prolonged conducted electrical weapon discharge in ethanol-intoxicated adults. *Am J Emerg Med* 2010;**28**:582–7.
77. Ho JD, Dawes DM, Bultman LL, Moscati RM, Janchar TA, Miner JR. Prolonged TASER use on exhausted humans does not worsen markers of acidosis. *Am J Emerg Med* 2009;**27**:413–8.
78. Ho JD, Dawes DM, Cole JB, Hottinger JC, Overton KG, Miner JR. Lactate and pH evaluation in exhausted humans with prolonged TASER X26 exposure or continued exertion. *Forensic Sci Int* 2009;**190**:80–6.
79. Dawes DM, Ho JD, Reardon RF, Miner JR. The cardiovascular, respiratory, and metabolic effects of a long duration electronic control device exposure in human volunteers. *Forensic Sci Med Pathol* 2010;**6**:268–74.
80. Eken C. Acidosis is a life-threatening condition regardless of the underlying condition. *Am J Emerg Med* 2008;**26**:721 [letter].
81. Byard RW, Wick R, Gilbert JD. Conditions and circumstances predisposing to death from positional asphyxia in adults. *J Forensic Leg Med* 2008;**15**:415–9.
82. Takeichi S, Wakasugi C, Shikata I. Fluidity of cadaveric blood after sudden death: Part I. Postmortem fibrinolysis and plasma catecholamine level. *Am J Forensic Med Pathol* 1984;**5**:223–7.
83. Hirvonen J, Kortelainen ML, Huttunen P. Pulmonary and serum surfactant phospholipids and serum catecholamines in strangulation. An experimental study on rats. *Forensic Sci Int* 1997;**90**:17–24.
84. Barton M, Lake CR, Rainey TG, Chernow B. Is catecholamine release pH mediated? *Crit Care Med* 1982;**10**:751–3.
85. Lim EC, Seet RC. TASER usage and neurological sequelae. *J Emerg Med* 2009;**37**:170–1 [letter].
86. Bui ET, Sourkes M, Wennberg R. Generalized tonic-clonic seizure after a taser shot to the head. *CMAJ* 2009;**180**:625–6.
87. Eastman AL, Metzger JC, Pepe PE, Benitez FL, Decker J, Rinnert KJ, et al. Conductive electrical devices: a prospective, population-based study of the medical safety of law enforcement use. *J Trauma* 2008;**64**:1567–72.
88. Mash DC. Biochemical brain markers in excited delirium deaths. In: Kroll MW, Ho JD, editors. *TASER® conducted electrical weapons: physiology, pathology, and law*. New York: Springer Science+Business Media; 2009. p. 365–77.
89. Bunai Y, Akaza K, Jiang WX, Nagai A. Fatal hyperthermia associated with excited delirium during an arrest. *Leg Med (Tokyo)* 2008;**10**:306–9.
90. Menaker J, Farcy DA, Boswell SA, Stein DM, Dutton RP, Hess JR, et al. Cocaine-induced agitated delirium with associated hyperthermia: a case report. *J Emerg Med*; in press. doi:10.1016/j.jemermed.2008.02.055.
91. Tsokos M. Postmortem changes and artifacts occurring during the early postmortem interval. Chapter 5. In: Tsokos M, editor. *Forensic pathology Reviews. Volume 3*. Totowa, New Jersey, USA: Humana Press Inc; 2006. p. 183–237.
92. Jauchem JR, Sherry CJ, Fines DA, Cook MC. Acidosis, lactate, electrolytes, muscle enzymes, and other factors in the blood of *Sus scrofa* following repeated TASER® exposures. *Forensic Sci Int* 2006;**161**:20–30.
93. Jauchem JR. Increased hematocrit after applications of conducted energy weapons (including TASER® devices) to *sus scrofa*. *J Forensic Sci* 2011;**56**(Suppl 1):S229–33.
94. Brun J-F, Varlet-Marie E, Connes P, Aloulou I. Hemorheological alterations related to training and overtraining. *Biorheology* 2010;**47**:95–115 [review].
95. Dawes DM, Ho JD, Reardon RF, Strote SR, Nelson RS, Lundin EJ, et al. The respiratory, metabolic, and neuroendocrine effects of a new generation electronic control device. *Forensic Sci Int*; in press. doi:10.1016/j.forsciint.2010.08.028.
96. Penttilä A, Laiho K. Autolytic changes in blood cells of human cadavers. II. Morphological studies. *Forensic Sci Int* 1981;**17**:121–32.
97. Cadroy Y, Pillard F, Sakariassen KS, Thalamas C, Boneu B, Riviere D. Strenuous but not moderate exercise increases the thrombotic tendency in healthy sedentary male volunteers. *J Appl Physiol* 2002;**93**:829–33.
98. Ikarugi H, Shibata M, Shibata S, Ishii H, Taka T, Yamamoto J. High intensity exercise enhances platelet reactivity to shear stress and coagulation during and after exercise. *Pathophysiol Haemost Thromb* 2003;**33**:127–33.
99. Ribeiro J, Almeida-Dias A, Oliveira AR, Mota J, Appell HJ, Duarte JA. Exhaustive exercise with high eccentric components induces prothrombotic and hypofibrinolytic responses in boys. *Int J Sports Med* 2007;**28**:193–6.
100. Kaufman MJ, Siegel AJ, Mendelson JH, Rose SL, Kukes TJ, Sholar MB, et al. Cocaine administration induces human splenic constriction and altered hematologic parameters. *J Appl Physiol* 1998;**85**:1877–83.
101. Jepson JH. Polycythemia: diagnosis, pathophysiology and therapy. II. *Can Med Assoc J* 1969;**100**:327–34 [review].
102. Nolte KB. Rhabdomyolysis associated with cocaine abuse. *Hum Pathol* 1991;**22**:1141–5.
103. Roth D, Alarcón FJ, Fernandez JA, Preston RA, Bourgoignie JJ. Acute rhabdomyolysis associated with cocaine intoxication. *N Engl J Med* 1988;**319**:673–7.
104. Asadollahi K, Hastings IM, Beeching NJ, Gill GV. Laboratory risk factors for hospital mortality in acutely admitted patients. *QJM* 2007;**100**:501–7.
105. Kreutziger J, Wenzel V, Kurz A, Constantinescu MA. Admission blood glucose is an independent predictive factor for hospital mortality in polytraumatized patients. *Intensive Care Med* 2009;**35**:1234–9.
106. Babu N, Singh M. Influence of hyperglycemia on aggregation, deformability and shape parameters of erythrocytes. *Clin Hemorheol Microcirc* 2004;**31**:273–80.
107. Lieber CS. Ethanol metabolism, cirrhosis and alcoholism. *Clin Chim Acta* 1997;**257**:59–84.
108. Vuong C, Van Uum SH, O'Dell LE, Lutfy K, Friedman TC. The effects of opioids and opioid analogs on animal and human endocrine systems. *Endocr Rev* 2010;**31**:98–132.
109. Molina DK, Hargrove VM. Fatal cocaine interactions: a review of cocaine-related deaths in Bexar County, Texas. *Am J Forensic Med Pathol* 2010; in press. doi:10.1097/PAF.0b013e3181ed79fe.
110. Mets B, Jamdar S, Landry D. The role of catecholamines in cocaine toxicity: a model for cocaine sudden death. *Life Sci* 1996;**59**:2021–31.
111. Pilgrim JL, Gerostamoulos D, Drummer OH, Bollmann M. Involvement of amphetamines in sudden and unexpected death. *J Forensic Sci* 2009;**54**:478–85.
112. Karch SB, Stephens BG, Ho CH. Methamphetamine-related deaths in San Francisco: demographic, pathologic, and toxicologic profiles. *J Forensic Sci* 1999;**44**:359–68.
113. Paterson DJ. Antiarrhythmic mechanisms during exercise. *J Appl Physiol* 1996;**80**:1853–62 [review].
114. O'Neill M, Paterson DJ. Role of the sympathetic nervous system in cardiac performance during hyperkalemia in the anesthetized pig. *Acta Physiol Scand* 1995;**153**:1–11.
115. Paterson DJ, Rogers J, Powell T, Brown HF. Effect of catecholamines on the ventricular myocyte action potential in raised extracellular potassium. *Acta Physiol Scand* 1993;**148**:177–86.
116. Dawes D, Ho J, Miner J. The neuroendocrine effects of the TASER® X26: a brief report. *Forensic Sci Int* 2009;**183**:14–9.
117. DiMaio V. Deaths in custody investigations. Chapter 10. In: Ross DL, Chan TC, editors. *Sudden deaths in custody*. 1st ed. Totowa, New Jersey: Humana Press; 2006. p. 167–72.
118. DiMaio TG, DiMaio VJM. *Excited delirium syndrome: cause of death and prevention*. Boca Raton, Florida: CRC/Taylor & Francis; 2006.
119. Ross DL, Chan TC. *Sudden deaths in custody*. 1st ed. Totowa, New Jersey: Humana Press; 2006.
120. Zhu BL, Ishikawa T, Michiue T, Li DR, Zhao D, Quan L, et al. Postmortem serum catecholamine levels in relation to the cause of death. *Forensic Sci Int* 2007;**173**:122–9.
121. Kernbach-Wighton G, Friedrich E, Rissler D, Niedmann PD, Saternus K-S. Post-mortem biochemical estimations in cases of lethal high excitation. *Rom J Leg Med* 2006;**14**:251–61.
122. Howlett K, Galbo H, Lorentsen J, Bergeron R, Zimmerman-Belsing T, Bülow J, et al. Effect of adrenaline on glucose kinetics during exercise in adrenalectomized humans. *J Physiol* 1999;**519**(Pt 3):911–21.
123. Zilg B, Alkass K, Berg S, Druid H. Postmortem identification of hyperglycemia. *Forensic Sci Int* 2009;**185**:89–95.
124. Madea B, Musshoff F. Postmortem biochemistry. *Forensic Sci Int* 2007;**165**:165–71 [review].
125. Glatter KA, Chiamvimonvat N, He Y, Chevalier P, Turillazzi E. Postmortem analysis for inherited ion channelopathies. Chapter 2. In: Rutty GN, editor. *Essentials of autopsy practice*. London, UK: Springer-Verlag; 2006. p. 15–37.
126. Tester DJ, Ackerman MJ. The role of molecular autopsy in unexplained sudden cardiac death. *Curr Opin Cardiol* 2006;**21**:166–72.
127. Kauerstein S, Kiehne N, Neumann T, Pitschner HF, Bratzke H. Cardiac gene defects can cause sudden cardiac death in young people. *Dtsch Arztebl Int* 2009;**106**:41–7 [review].
128. Cerretani D, Riezzo I, Fiaschi AI, Centini F, Giorgi G, D'Errico S, et al. Cardiac oxidative stress determination and myocardial morphology after a single ecstasy (MDMA) administration in a rat model. *Int J Leg Med* 2008;**122**:461–9.
129. Hua YS, Liang R, Liang L, Huang GZ. Contraction band necrosis in two ecstasy abusers: a latent lethal lesion associated with ecstasy. *Am J Forensic Med Pathol* 2009;**30**:295–7.

130. Mash DC, Duque L, Pablo J, Qin Y, Adi N, Hearn WL, et al. Brain biomarkers for identifying excited delirium as a cause of sudden death. *Forensic Sci Int* 2009;**190**:e13–9.
131. Maeda H, Zhu BL, Bessho Y, Ishikawa T, Quan L, Michiue T, et al. Postmortem serum nitrogen compounds and C-reactive protein levels with special regard to investigation of fatal hyperthermia. *Forensic Sci Med Pathol* 2008;**4**:175–80.
132. Nixdorf-Miller A, Hunsaker DM, Hunsaker 3rd JC. Hypothermia and hyperthermia medicolegal investigation of morbidity and mortality from exposure to environmental temperature extremes. *Arch Pathol Lab Med* 2006;**130**:1297–304 [review].
133. Zhu BL, Ishikawa T, Michiue T, Li DR, Zhao D, Oritani S, et al. Postmortem cardiac troponin T levels in the blood and pericardial fluid. Part 1. Analysis with special regard to traumatic causes of death. *Leg Med (Tokyo)* 2006;**8**:86–93.
134. Maeda H, Michiue T, Zhu BL, Ishikawa T, Quan L. Analysis of cardiac troponins and creatine kinase MB in cerebrospinal fluid in medicolegal autopsy cases. *Leg Med (Tokyo)* 2009;**11**(Suppl. 1):S266–8.
135. Zhu BL, Ishikawa T, Michiue T, Li DR, Zhao D, Quan L, et al. Evaluation of postmortem urea nitrogen, creatinine and uric acid levels in pericardial fluid in forensic autopsy. *Leg Med (Tokyo)* 2005;**7**:287–92.
136. Li DR, Quan L, Zhu BL, Ishikawa T, Michiue T, Zhao D, et al. Evaluation of postmortem calcium and magnesium levels in the pericardial fluid with regard to the cause of death in medicolegal autopsy. *Leg Med (Tokyo)* 2009;**11**(Suppl. 1):S276–8.
137. Ishikawa T, Quan L, Li DR, Zhao D, Michiue T, Hamel M, et al. Post-mortem biochemistry and immunohistochemistry of adrenocorticotrophic hormone with special regard to fatal hypothermia. *Forensic Sci Int* 2008;**179**:147–51.
138. Quan L, Ishikawa T, Michiue T, Li DR, Zhao D, Zhu BL, et al. Quantitative analysis of ubiquitin-immunoreactivity in the midbrain periaqueductal gray matter with regard to the causes of death in forensic autopsy. *Leg Med (Tokyo)* 2005;**7**:151–6.
139. Nishio H, Suzuki K. Three cases of suspected hyperthermia with remarkable elevation of serum mast cell tryptase. *Forensic Sci Int* 2005;**149**:51–5.
140. Yoshida C, Ishikawa T, Michiue T, Quan L, Maeda H. Postmortem biochemistry and immunohistochemistry of chromogranin A as a stress marker with special regard to fatal hypothermia and hyperthermia. *Int J Leg Med* 2011;**125**:11–20.
141. Ishikawa T, Yoshida C, Michiue T, Perdekamp MG, Pollak S, Maeda H. Immunohistochemistry of catecholamines in the hypothalamic-pituitary-adrenal system with special regard to fatal hypothermia and hyperthermia. *Leg Med (Tokyo)* 2010;**12**:121–7.
142. Payne-James J. History and development of clinical forensic medicine. Chapter 1. In: Stark MM, editor. *Clinical forensic medicine: a physician's guide*. Totowa, New Jersey, USA: Humana Press Inc.; 2005. p. 1–36.
143. Hall CA. [Les pistolets Taser, un risque inacceptablement élevé ou suffisamment bas pour être acceptable?] Taser guns, an unacceptably high risk or sufficiently low to be acceptable? *CJEM* 2009;**11**:84–6. 87–9 [French, English] [editorial].
144. Kroll MW, Panescu D, Brewer JE, Lakkireddy DJ, Graham M. *Weight-adjusted meta-analysis of fibrillation risk from TASER® conducted electrical weapons*. American Academy of Forensic Sciences. Proceedings of the 61st Annual Meeting ("Forensic science: envisioning and creating the future"), Denver, Colorado; February 16–21, 2009. pp. 177–8.
145. Southall P, Grant J, Fowler D, Scott S. Police custody deaths in Maryland, USA: an examination of 45 cases. *J Forensic Leg Med* 2008;**15**:227–30.
146. Bozeman WP, Hauda 2nd WE, Heck JJ, Graham Jr DD, Martin BP, Winslow JE. Safety and injury profile of conducted electrical weapons used by law enforcement officers against criminal suspects. *Ann Emerg Med* 2009;**53**:480–9.
147. *Calgary Police Commission. Minutes of open meeting*. Calgary, Alberta, Canada. Available at http://www.calgary.ca/docgallery/bu/cpc/meetings/2006/september_minutes.pdf; September 19, 2006.
148. Bozeman WP, Winslow JE. Medical aspects of less lethal weapons. *Internet J Rescue Disaster Med* (1), <http://www.ispub.com/ostia/index.php?xmlFilePath=journals/ijrdm/vol5n1/lethal.xml>, 2005;**5**.
149. Glick J. 'Excited delirium' cited in deaths. *Police One*, <http://www.policeone.com/edp/articles/1179873-Excited-Delirium-cited-in-deaths/>, September 2006;**25**.
150. Ho JD. Some more food for thought on the current TASER device literature. *Am Coll Emerg Physicians Web Page*, <http://www.acep.org/content.aspx?id=31840>; 2008.
151. Young JG, Chiasson DA, Cairns JT, Carter EJ, Payne-James J. Custody and restraint deaths. In: Payne-James J, Busuttill A, Smock W, editors. *Forensic medicine: clinical and pathological aspects*. San Francisco, London: Greenwich Medical Media Ltd; 2003. p. 115–31.
152. Grant JR, Southall PE, Mealey J, Scott SR, Fowler DR. Excited delirium deaths in custody: past and present. *Am J Forensic Med Pathol* 2009;**30**:1–5.
153. Stratton SJ, Rogers C, Brickett K, Gruzinski G. Factors associated with sudden death of individuals requiring restraint for excited delirium. *Am J Emerg Med* 2001;**19**:187–91.
154. Sathyavagiswaran L, Rogers C, Noguchi TT. Restraint asphyxia in in-custody deaths Medical examiner's role in prevention of deaths. *Leg Med (Tokyo)* 2007;**9**:88–93.
155. Cina SJ, Davis GJ. Cardiac dysrhythmia during restraint. *Am J Forensic Med Pathol* 2010;**31**:e4 [letter].